

# On the (Im-)Possibility of Representing Probability Distributions as a Difference of I.I.D. Noise Terms

Christian Ewerhart,<sup>a,\*</sup> Marco Serena<sup>b</sup>

<sup>a</sup>Department of Economics, University of Zurich, 8001 Zurich, Switzerland; <sup>b</sup>School of Business, Economics and Law, CUNEF Universidad, 28040 Madrid, Spain

\*Corresponding author

Contact: christian.ewerhart@econ.uzh.ch,  <https://orcid.org/0000-0003-3285-1099> (CE); marco.serena@cunef.edu,  <https://orcid.org/0000-0002-6215-9629> (MS)

Received: March 16, 2023

Revised: October 16, 2023

Accepted: January 7, 2024

Published Online in Articles in Advance:  
March 7, 2024

MSC2020 Subject Classifications: Primary:  
60E05; Secondary: 91B26

<https://doi.org/10.1287/moor.2023.0081>

Copyright: © 2024 INFORMS

**Abstract.** A random variable is *difference-form decomposable* (DFD) if it may be written as the difference of two i.i.d. random terms. We show that densities of such variables exhibit a remarkable degree of structure. Specifically, a DFD density can be neither approximately uniform, nor quasiconvex, nor strictly concave. On the other hand, a DFD density need, in general, be neither unimodal nor logconcave. Regarding smoothness, we show that a compactly supported DFD density cannot be analytic and will often exhibit a kink even if its components are smooth. The analysis highlights the risks for model consistency resulting from the strategy widely adopted in the economics literature of imposing assumptions directly on a difference of noise terms rather than on its components.

**Keywords:** differences of random variables • density functions • characteristic function • uniform distribution

## 1. Preliminaries

### 1.1. Introduction

In a large variety of economic models, uncertainty enters the framework in the form of the difference of two i.i.d. random variables, say  $\varepsilon_1$  and  $\varepsilon_2$ . For instance, in a rank-order tournament à la Lazear and Rosen [38], individual performance is the sum of input and some randomness, and the winner is determined by comparing levels of individual performance across contestants. The economic prediction (or power of statistical test) may then crucially depend on the distribution of the difference of the noise terms,  $\varepsilon_1 - \varepsilon_2$ . This is similarly the case in a variety of other settings, including contests (Che and Gale [14], Hirshleifer [33]), models of location choice (Moretti [49], Roback [59], Rosen [60]), vertical differentiation (Lin [40]), probabilistic voting (Glaeser et al. [29], Lindbeck and Weibull [41]), random utility (Becker et al. [8], Goeree et al. [30]), and paired comparisons (Bradley and Terry [10], Thurstone [66]). For the applied economist entrusted with the analysis of such models, it may appear natural and convenient to impose assumptions directly on the distribution of the difference rather than on the distribution of the components  $\varepsilon_1$  and  $\varepsilon_2$ . This approach, however, is not entirely innocuous. Indeed, as will be discussed, certain familiar probability distributions, such as the uniform distribution, simply cannot be represented as the difference of two i.i.d. random variables. However, this fact is rarely mentioned and sometimes even neglected in economic modeling. Thus, there is the risk of ending up with an inconsistent set of assumptions. At the same time, there does not seem to be a single reference that offers help on this issue.<sup>1</sup>

The present paper aims at providing an initial, systematic, and accessible study of the class of random variables that correspond to the difference of two i.i.d. random variables. We refer to such random variables as *difference-form decomposable* (DFD). It turns out that density functions of DFD random variables exhibit a remarkable degree of structure. Starting from the important special case of the uniform distribution, the analysis identifies several broad classes of distributions that have the property of not being DFD. We show that random variables with an approximately uniform, quasiconvex, or strictly concave density function cannot be DFD. The observation that a strictly concave density never corresponds to a DFD random variable may be of particular interest. To prove this *necessary* condition, we recycle the argument underlying Pólya's [55] *sufficient* condition for a function to be the characteristic function of a real-valued random variable. To illustrate the applicability of this result, we note that the strictly concave beta density is never DFD. We also show that other properties that may be expected from a DFD density, such as unimodality or logconcavity, need not hold in general (but do so if the components possess these properties).

To study the smoothness properties of DFD densities, we analyze the limit behavior of characteristic functions using a theorem of Erdélyi [24]. Results are obtained for compactly supported densities. Specifically, it turns out that any continuously differentiable DFD density necessarily vanishes at the lower and upper boundaries of its support interval. A similar relationship holds for the higher derivatives. Going to the limit, we find that a compactly supported DFD density is never analytic; that is, there must be at least one point in its support interval where the density cannot be approximated arbitrarily well by its Taylor series. This necessary condition is easy to apply. For example, we use it to show that neither the beta distribution with integer parameter nor the raised cosine distribution can be DFD. We also point out that a DFD density will often exhibit kinks even if its difference-form components are smooth.

We go on and explore sufficient conditions for difference-form decomposability. We note that any random variable with the property that the square root of its characteristic function is a *positive definite* function is DFD. An example is the class of infinitely divisible distributions. This sufficient condition is strong enough to allow the computation of the (identical) densities of the components  $\varepsilon_1$  and  $\varepsilon_2$  from a given distribution of the difference  $\varepsilon_1 - \varepsilon_2$ . Several extensions are offered as well. Specifically, we discuss distributions with finite support, functional inequalities (i.e., inequalities that restrict the values of DFD densities), ratio-form decomposability, and correlated noise terms. The extension regarding ratio-form decomposability sheds light, in particular, on Jia's [37] characterization of the Tullock contest technology.

For the reader's convenience, the results previewed above will be formulated without reference to the complex numbers. In particular, the characteristic function is generally expressed as a cosine transform, which is an ordinary integral of a real-valued integrand. As we believe, this helps intuition but also simplifies the application of our results.

## 1.2. Economic Motivation

Despite Lazear and Rosen [38] having noted the impossibility of an i.i.d. difference being uniform,<sup>2</sup> this fact is rarely discussed in the literature. In fact, screening the economics literature on the before-mentioned applications where difference-form decomposability emerges quite naturally, we found numerous papers assuming uniform differences. Very few of those papers, however, are explicit about the problem.

Working with an inconsistent set of assumptions is risky not only because the conclusions may become shaky but also because the intuitive interpretation of the model may become difficult. In a standard tournament, for instance, assuming that  $\varepsilon_1 - \varepsilon_2$  is uniform would render the marginal probability of winning independent of the opponent's effort, and therefore blur the borderline between relative performance evaluation and individual contracts.<sup>3</sup> Similarly, in a model of location choice or in a random utility model, assuming a non-DFD stochastic difference of the utility of two options would be at odds with the usual understanding that individual choices have a well-defined utility level after the resolution of uncertainty. Finally, a statistical test might be biased if the difference of two noise terms is assumed to follow a distribution that is not DFD.

In those and other economic models, the question of whether a density is DFD naturally emerges. It thus seems important to understand which random variables can be represented as the difference of two i.i.d. terms, and how stochastic properties of the components relate to properties of the difference term.

## 1.3. Related Literature

We are not aware of any work that tried characterizing the set of DFD distributions. Notwithstanding, almost any treatment of the topic of characteristic functions, our main tool of investigation, touches upon the matter (see, e.g., Linnik [42] or Lukacs [44]). Feller [26] explains how *symmetrization* helps avoid messy arguments in probability theory, for example, because the tails of the distribution functions of a random variable and its symmetrization are of comparable magnitude. The celebrated 123 Theorem by Alon and Yuster [3] offers inequalities satisfied by DFD distribution functions and is, therefore, complementary to our investigation of DFD densities.

The indecomposability of the continuous uniform distribution dates back to Puri and Sen [58, p. 970], who thank Basu and Puri for the short proof. We have not seen attempts to generalize this result, however. Our criterion that strict concavity of the density is incompatible with being DFD is related to Pólya's [55] sufficient condition for being a characteristic function (see also Tuck [69]).

In optics and crystallography, the task of recovering a measure from its symmetrization or, equivalently, from the modulus of its characteristic function is known as *phase retrieval*. This problem arises in optics because the measurement of a diffracted wavefront (e.g., resulting from a beam of laser light sent through a gap) gives only the intensity of the wave form rather than its complex amplitude (Patterson [54], Rosenblatt [61], Walther [72]). In related work, Giraud and Peschanski [28] and Gori [31] studied nonnegative functions whose Fourier transform is likewise nonnegative.<sup>4</sup>

## 1.4. Overview of the Paper

The remainder of this paper is structured as follows. In Section 2, we introduce the notion of difference-form decomposability. Section 3 deals with conditions on the shape of DFD densities. Section 4 discusses smoothness conditions. Section 5 derives sufficient conditions for difference-form decomposability and a formula for the construction of the difference-form component. Section 6 offers extensions. Section 7 concludes. Technical proofs have been relegated to an appendix.

## 2. Difference-Form Decomposability

In this section, we introduce the class of DFD distributions. We will provide basic definitions (Section 2.1), review the necessary background on characteristic functions (Section 2.2), survey examples of DFD distributions (Section 2.3), and similarly survey examples of distributions that are not DFD (Section 2.4).

### 2.1. Basic Definitions

All random variables considered in this paper are assumed to be real-valued. The following concept is central to our analysis.

**Definition 1.** Let  $Z$  be a random variable. We will say that  $Z$  is **difference-form decomposable (DFD)** if there are two i.i.d. random variables  $X$  and  $Y$  such that  $Z \stackrel{d}{=} X - Y$ .

The equation  $Z \stackrel{d}{=} X - Y$  says that  $Z$  and  $X - Y$  follow the same probability law. When a random variable  $Z$  is DFD as specified in Definition 1, then the random variable  $X$  is referred to as a *difference-form component* of  $Z$ .<sup>5</sup>

To understand the specific nature of the analysis pursued in the present paper, it is important to acknowledge that the components  $X$  and  $Y$  are required to be *identically* distributed. Indeed, dropping that requirement would lead to an entirely different research question, easily seen to be equivalent to the problem of additive decomposability. That problem, however, has been well studied (e.g., Linnik [42], Linnik and Ostrovskiĭ [43]). Similarly important is the requirement that  $X$  and  $Y$  are *independent* (see also Section 6). For example, if  $X$  and  $Y$  are uniform and perfectly negatively correlated, then  $Z$  is again uniform (Bagnoli et al. [6], Meyer [47]). Thus, the i.i.d. requirement is crucial to all that follows.<sup>6</sup>

**Definition 2.** A random variable  $Z$  is called **symmetric** (about zero) if  $Z \stackrel{d}{=} -Z$ .

The following observation is simple but important.

**Lemma 1.** Any DFD random variable  $Z$  is symmetric.

**Proof.** Suppose that  $Z$  is DFD. Then, there exist i.i.d. random variables  $X$  and  $Y$  such that  $Z \stackrel{d}{=} X - Y$ . But then,  $-Z \stackrel{d}{=} Y - X$ , which proves the claim.  $\square$

In view of Lemma 1, a DFD random-variable  $Z$  with component  $X$  is commonly known as the *symmetrization* of  $X$ .

Suppose that  $X$  is a difference-form component of  $Z$ . Then the distribution functions of  $Z$  and  $X$  will be denoted by  $G = G(z)$  and  $F = F(x)$ , respectively. In view of our applications, we will mostly focus on continuous distributions, that is, on distributions that admit a density. If densities exist, these will correspondingly be denoted by  $g = g(z)$  and  $f = f(x)$ . Moreover, the convolution relationship

$$g(z) = \int_{-\infty}^{\infty} f(x+z)f(x)dx$$

admits an interpretation as an autocorrelation function. To avoid clumsy language, the concepts introduced above, that is, DFD, difference-form component, and symmetry, will be informally extended to distributions and density functions.

### 2.2. Characteristic Functions

In general, the *characteristic function* of a random variable  $Z$  is defined as  $\varphi_Z(t) = E[e^{itZ}] \equiv \int_{-\infty}^{\infty} e^{itz} dG(z)$ , where  $i = \sqrt{-1}$ , and the parameter  $t$  is real (e.g., Lukacs [44]). The relevance of the characteristic function for the problem at hand is the following important observation.

**Lemma 2.** Suppose that  $Z$  is DFD with component  $X$ . Then,  $\varphi_Z(t) = |\varphi_X(t)|^2 \geq 0$  for all  $t \in \mathbb{R}$ , where  $\varphi_X(t)$  denotes the characteristic function of  $X$ .

**Proof.** The characteristic function of  $-X$  is given by  $\varphi_{-X}(t) = E[e^{-itX}]$ . Because  $t \in \mathbb{R}$  and  $X$  is real-valued, this implies  $\varphi_{-X}(t) = \overline{\varphi_X(t)}$ , where the upper bar denotes complex conjugation. Hence, by the multiplication theorem for characteristic functions,  $\varphi_Z(t) = \varphi_X(t)\varphi_{-X}(t) = \varphi_X(t)\overline{\varphi_X(t)} = |\varphi_X(t)|^2$ . In particular,  $\varphi_Z(t) \geq 0$  for any  $t \in \mathbb{R}$ . The claim follows.  $\square$

Thus, a necessary condition for  $Z$  to be DFD is that its characteristic function is real-valued and nonnegative. Using Lemma 2 as a first numerical check of decomposability of a given density is feasible. However, the reader is cautioned that the computation of Fourier transforms is not trivial (e.g., Ahmed et al. [1]). Moreover, the nonnegativity condition on the integral transform is not sufficient for  $Z$  to be DFD. A counterexample due to Lukacs [44] is replicated below. In fact, we will exhibit a more elementary counterexample in our discussion of distributions with finite support.

Although the characteristic function is, in general, complex-valued, there is an intuitive representation of  $\varphi_Z(t)$  as a cosine transform if  $Z$  is symmetric (e.g., DFD).

**Lemma 3.** If  $Z$  is symmetric, then  $\varphi_Z(t) = \int_{-\infty}^{\infty} \cos(tz) dG(z)$ , for any  $t \in \mathbb{R}$ .

**Proof.** Because  $Z$  is symmetric,  $E[e^{itZ}] = \frac{1}{2}E[e^{itZ} + e^{-itZ}] = E[\cos(tZ)]$ .  $\square$

The advantage of expressing the characteristic function as a cosine transform is that no reference to complex numbers is needed. In addition, there is an intuitive interpretation now that is not so immediate otherwise. Specifically, the cosine factor works like an *amplitude modulation* (AM). Amplitude modulation is used in electronic communication, radio transmission, computer modems, etc., to transmit a low-frequency audio signal via a high-frequency radio signal (see, e.g., Carson [13]). Similarly, the density of  $Z$  may be understood to modulate the cosine signal of a given frequency  $t$ , so that some information about the distribution of  $Z$  is captured in  $\varphi_Z(t)$ .<sup>7</sup>

### 2.3. Examples of Distributions That Are DFD

Table 1 provides details on the difference-form decomposability of several standard families of probability distributions. See also Figure 1 for an illustration of the corresponding density functions. The list starts with the examples of the normal and Cauchy distributions, both of which are infinitely divisible. As these examples suggest, any infinitely divisible distribution that is symmetric about zero is DFD. Particularly strong implications are feasible for normal distributions. Indeed, by Cramér's [16] theorem, any nondegenerate component of a normal distribution is normal. Thus, the difference-form decomposition shown in Table 1 is essentially unique in the normal case. Conversely, as shown by Carnal and Dozzi [12], this uniqueness property regarding difference-form decomposability is shared by no other infinitely divisible distribution. In the general case (i.e., if the distribution is not necessarily infinitely divisible), one can still show that any representation of a random variable as the difference of two *symmetric* i.i.d. noise terms is essentially unique. However, the difference-form decomposition is not unique in general if one allows for components that are not symmetric. For instance, the Laplace distribution may be represented either as the difference of two symmetric Bessel distributions or as the difference of two exponential distributions (which are not symmetric).

An example of a DFD distribution with compact support is the *triangular* distribution. Apart from this example, we do not know of any DFD distribution with compact support that has been (correctly) used in applications.<sup>8</sup>

### 2.4. Examples of Distributions That Are Not DFD

Table 2 lists a number of distributions that are not DFD. Particularly prominent is the example of the uniform distribution. But as will be shown below, the beta distribution (under parameter constraints) and the raised cosine

**Table 1.** Examples of difference-form decomposable distributions.

Distribution, parameter	Support	Density function $g(z)$	Characteristic function $\varphi_Z(t)$	Component	Support	Density function $f(x)$	Characteristic function $\varphi_X(t)$
Normal ( $\sigma_Z > 0$ )	$\mathbb{R}$	$\frac{1}{\sqrt{2\pi}\sigma_Z} \exp\left(-\frac{z^2}{2\sigma_Z^2}\right)$	$\exp\left(-\frac{\sigma_Z^2 t^2}{2}\right)$	Normal ( $\sigma_X^2 = \sigma_Z^2/2$ )	$\mathbb{R}$	$\frac{1}{\sqrt{2\pi}\sigma_X} \exp\left(-\frac{x^2}{2\sigma_X^2}\right)$	$\exp\left(-\frac{\sigma_X^2 t^2}{2}\right)$
Cauchy ( $\gamma_Z > 0$ )	$\mathbb{R}$	$\frac{\gamma_Z}{\pi(\gamma_Z^2 + z^2)}$	$\exp(-\gamma_Z  t )$	Cauchy ( $\gamma_X = \gamma_Z/2$ )	$\mathbb{R}$	$\frac{\gamma_X}{\pi(\gamma_X^2 + x^2)}$	$\exp(-\gamma_X  t )$
Logistic ( $\beta > 0$ )	$\mathbb{R}$	$\frac{\exp(-z/\beta)}{\beta(1+\exp(-z/\beta))^2}$	$\frac{\pi\beta t}{\sinh(\pi\beta t)}$	Gumbel	$\mathbb{R}$	$\frac{1}{\beta} \exp\left(-\frac{x}{\beta} - \exp\left(-\frac{x}{\beta}\right)\right)$	$\Gamma(1 - i\beta t)$
Laplace ( $\lambda > 0$ )	$\mathbb{R}$	$\frac{\lambda}{2} \exp(-\lambda  z )$	$\frac{\lambda^2}{\lambda^2 + t^2}$	Bessel	$\mathbb{R}$	$\frac{\lambda}{\pi} K_0(\lambda  x )$	$\frac{\lambda}{\sqrt{\lambda^2 + t^2}}$
				Exponential	$\mathbb{R}_{\geq 0}$	$\lambda \exp(-\lambda x)$	$\frac{\lambda}{\lambda - it}$
Triangular ( $c > 0$ )	$[-2c, 2c]$	$\max\left\{\frac{1}{2c} - \frac{ z }{4c^2}, 0\right\}$	$\frac{\sin^2(ct)}{c^2 t^2}$	Uniform	$[-c, c]$	$\begin{cases} 1/(2c) & \text{if } x \in [-c, c] \\ 0 & \text{otherwise} \end{cases}$	$\frac{\sin(ct)}{ct}$

**Table 2.** Examples of distributions that are *not* difference-form decomposable.

Distribution, parameter	Support	Density function $g(z)$	Characteristic function $\varphi_Z(t)$	Reference
Uniform ( $c > 0$ )	$[-c, c]$	$\begin{cases} 1/(2c) & \text{if } z \in [-c, c] \\ 0 & \text{otherwise} \end{cases}$	$\frac{\sin(ct)}{ct}$	Puri and Sen [57], this paper (Example 1)
$n/a$		Any approximately uniform	$n/a$	This paper (Proposition 1)
Beta ( $0 < \alpha \leq 2$ or $\alpha \in \mathbb{N}$ )	$[-1, 1]$	$\frac{\Gamma(\frac{\alpha+1}{2})}{\sqrt{\pi}\Gamma(\frac{\alpha}{2})}(1-z^2)^{\alpha-1}$	Confluent hypergeometric	This paper (Example 2)
$n/a$		Any quasiconvex	$n/a$	This paper (Proposition 2)
$n/a$		Any strictly concave	$n/a$	This paper (Theorem 1)
Raised cosine ( $c > 0$ )	$[-c, c]$	$\begin{cases} \frac{1 + \cos(z\pi/c)}{2c} & \text{if } z \in [-c, c] \\ 0 & \text{otherwise} \end{cases}$	$\frac{\pi^2 \sin(ct)}{ct(\pi^2 - c^2 t^2)}$	This paper (Example 3)
$n/a$		Any compactly supported analytic	$n/a$	This paper (Proposition 3)
$n/a$	$\mathbb{R}$	$\frac{1}{\sqrt{2\pi}} z^2 \exp\left(-\frac{z^2}{2}\right)$	$(1-t^2)\exp\left(-\frac{t^2}{2}\right)$	Dugué [20, p. 38], this paper (Lemma 2)
$n/a$	$\mathbb{R}$	$\sqrt{\frac{2}{3\pi}} (8z^2 - 1)^2 \exp\left(-\frac{2}{3}z^2\right)$	$\left(1 - \frac{t^2}{2}\right)^2 \exp\left(-\frac{3}{8}t^2\right)$	Lukacs [44, p. 127]


Note. Only distributions that are symmetric with respect to the origin are listed (cf. Lemma 1).

distribution likewise fall in this class. All these examples have compact support. Dugué [20] constructed a distribution with full support that is not DFD. Although that example is still covered by Lemma 2 above, the more complicated example of Lukacs [44] even has a nonnegative characteristic function, and therefore illustrates the fact that the conclusion of Lemma 2 is only necessary, but not sufficient, for a random variable to be DFD. In fact, both distributions are entirely indecomposable (i.e., even allowing for nontrivial heterogeneous factors). The density functions of these examples are illustrated in Figure 2.

### 3. Shape Conditions


This section starts the formal analysis by deriving a variety of necessary conditions on the shape of DFD distributions. We will discuss uniform and approximately uniform distributions (Section 3.1), quasiconvex densities (Section 3.2),

**Figure 1.** Examples of difference-form decomposable distributions.



Note. Shown are densities of the normal (solid), Cauchy (dashed), logistic (crossed), Laplace (thin), and triangular (dotted) distributions.



**Figure 2.** Examples of distributions that are not difference-form decomposable.

Note. Shown are the densities of the uniform (solid), raised cosine (dashed), concave beta (diamonds), convex beta (thin), Dugué (bold), and Lukacs (crosses) distributions.

and strictly concave densities (Section 3.3). The present section concludes with a discussion of unimodal and logconcave densities (Section 3.4).

### 3.1. Uniform and Approximately Uniform Distributions


The following classic result sets the stage for our analysis.

**Example 1.**  $Z \sim U[-1, 1]$  is not DFD.

This observation, a formal proof of which is included in the appendix, is stated in Puri and Sen [58, p. 970], for instance.<sup>9</sup> Example 1 also shows that symmetry is not sufficient for a random variable to be DFD. To understand how this observation follows from Lemma 2 (the nonnegativity of the cosine transform), see Figure 3. The point to note is that the signed area between the plotted graph and the horizontal axis is negative. Intuitively, *the uniform density declines too slowly*, so that the integral  $\varphi_Z(t)$  turns negative for a suitably chosen value of  $t$ , which is inconsistent with difference-form decomposability. For  $t = \frac{3\pi}{2}$ , for instance, one obtains

$$\varphi_Z(t) = \frac{1}{2} \int_{-1}^1 \cos\left(\frac{3\pi z}{2}\right) dz = -\frac{2}{3\pi} < 0.$$

This, however, is in conflict with the conclusion of Lemma 2.<sup>10</sup>

**Figure 3.** Plot of the function  $z \mapsto \cos\left(\frac{3}{2}\pi z\right)$  over the interval  $[0, 1]$ .

Below, we will see that broad classes of distributions that either contain the uniform distribution as a special case or feature it as a limit case are likewise not DFD. This will be interpreted as evidence showing that the lack of decomposability of the uniform distribution, which is particularly relevant for economic applications, is a robust phenomenon rather than a knife-edge pathology of limited relevance.

To start with, one might ask if, despite a uniform distribution not being DFD, there exist two i.i.d. random variables whose difference can at least *approximate* a uniform distribution. The answer is negative.

**Proposition 1.** *Any density function on the interval  $[-1, 1]$  that has values in the interval  $(\frac{1}{2} - \delta, \frac{1}{2} + \delta)$ , where  $\delta > 0$  is small, is not DFD.*

**Proof.** See the appendix.  $\square$

The proof shows that  $\delta = \frac{1}{6} \approx 0.166$ . To understand why Proposition 1 holds true, one takes another look at Figure 3 and notes that the area with the negative weight, where  $g(z) > \frac{1}{2} - \frac{1}{6} = \frac{1}{3}$ , is precisely twice as large as the area with the positive weight, where  $g(z) < \frac{1}{2} + \frac{1}{6} = \frac{2}{3}$ . Thus, again, the cosine transform has a negative sign.

### 3.2. Quasiconvex Densities

Although Proposition 1 is useful, it cannot deal with the following example.

**Example 2.** The density of the symmetric beta distribution is given as

$$g(z) = \frac{\Gamma(\alpha + \frac{1}{2})}{\sqrt{\pi}\Gamma(\alpha)} (1 - z^2)^{\alpha-1} \quad (z \in [-1, 1]),$$

where  $\alpha > 0$  is a shape parameter.<sup>11</sup> For  $\alpha \in (0, 1]$ , this density is convex, where the boundary case  $\alpha = 1$  corresponds to the uniform distribution. For  $\alpha \in (1, 2]$ , the density is strictly concave. For  $\alpha > 2$ , however,  $g$  is neither convex nor concave.

In the example, the conditions of Proposition 1 are not satisfied (unless  $\alpha = 1$ ). However, the intuition underlying the uniform case admits another generalization, namely, to the class of quasiconvex density functions. We say that a density function  $g$  of a symmetric distribution with compact support  $[-c, c]$ , where  $c > 0$ , is *quasiconvex* if and only if  $g$  is weakly decreasing on  $(-c, 0]$  and weakly increasing on  $[0, c)$  (i.e., we disregard the boundary points of the support for convenience).

**Proposition 2.** *A quasiconvex density cannot be DFD.*

**Proof.** See the appendix.  $\square$

Like Proposition 1 above, this result is derived by extending the graphical proof of the uniform case. If the symmetric density is weakly increasing on  $[0, 1)$ , then the signed integral outlined in Figure 3 is weakly smaller than for the uniform distribution, which is already negative.

Proposition 2 indeed allows dealing with additional parameter values in the example of the beta distribution. Specifically, if the density is convex (i.e., if  $\alpha \in (0, 1]$ ), then the beta distribution cannot be DFD.

Further below, we will obtain a variant of Proposition 2 saying that a random variable represented by a continuous density function  $g$  on  $[-1, 1]$  does not admit a difference-form component with continuously differentiable density unless  $g$  assumes its maximum at the origin.


### 3.3. Strictly Concave Densities

For  $\alpha \in (1, 2]$ , the density of the beta distribution is strictly concave. Hence, our previous criteria do not apply. This case is, however, covered by the following result, which for us was the least expected finding of the present analysis.

**Proposition 3.** *A density that is strictly concave on its support cannot be DFD.*

**Proof.** See the appendix.  $\square$

The proof of Proposition 3 is inspired by Pólya's [55] sufficient condition for characteristic functions.<sup>12</sup> The first point to note is that a strictly concave density must be compactly supported, say on  $[-1, 1]$ . Next, one notes that the evaluation of the cosine transform at the special value  $t = 2\pi$  decomposes the interval  $[0, 1]$  into four subintervals of length  $\frac{1}{4}$ . Moreover, one may easily check that  $\cos(2\pi z) = -\cos(2\pi(\frac{1}{2} - z)) = -\cos(2\pi(\frac{1}{2} + z)) = \cos(2\pi(1 - z))$ , for any  $z \in [0, \frac{1}{4}]$ . See Figure 4 for an illustration of this relationship. Hence, the integrand of the cosine transform

**Figure 4.** Intuition underlying the proof of Proposition 3.

satisfies

$$\begin{aligned}
 & g(z)\cos(2\pi z) + g\left(\frac{1}{2} - z\right)\cos\left(2\pi\left(\frac{1}{2} - z\right)\right) \\
 & + g\left(\frac{1}{2} + z\right)\cos\left(2\pi\left(\frac{1}{2} + z\right)\right) + g(1 - z)\cos(2\pi(1 - z)) \\
 & = \underbrace{\left(g(z) - g\left(\frac{1}{2} - z\right) - g\left(\frac{1}{2} + z\right) + g(1 - z)\right)}_{<0} \underbrace{\cos(2\pi z)}_{>0} < 0,
 \end{aligned}$$

as a consequence of strict concavity of  $g$ . Integrating over  $[0, \frac{1}{4}]$ , the cosine transform is seen to be negative, that is,  $\varphi_Z(2\pi) < 0$ , in conflict with Lemma 2. The proof given in the appendix works with partial integration like Pólya's original proof but captures the very same intuition.

### 3.4. Unimodal and Logconcave Densities

Unimodal densities are widely used in economics. It is known that the difference (but not necessarily the sum; see Chung [15]) of two i.i.d. unimodal random variables is necessarily unimodal (Hodges and Lehmann [34], Vogt [71]; see also Barlevy and Neal [7], for application). The same is true for strongly unimodal (i.e., logconcave) densities (An [5], Ibragimov [36]). As mentioned before, we will later show that any DFD density with well-behaved components necessarily assumes its maximum at the origin. However, even though unimodality with mode at zero seems to be a common feature of many DFD densities, neither unimodality nor logconcavity is necessary for a density to be DFD. For example, let  $f(x) = \frac{3}{2}$  if  $x \in [0, \frac{1}{3}] \cup [\frac{2}{3}, 1]$ , and  $f(x) = 0$  otherwise. Clearly,  $f$  is a density. Let  $X, Y$  be i.i.d. according to  $f$ . Then, the random variable  $Z = X - Y$  does not admit a unimodal density  $g$ . Unimodality and, similarly, logconcavity are neither sufficient for, say, a symmetric density to be DFD, as follows immediately from Proposition 3.

## 4. Smoothness

In this section, we will derive necessary conditions that rely on smoothness properties of the DFD density. We first discuss boundary conditions (Section 4.1), then analyticity (Section 4.2), and finally kinks (Section 4.3).

### 4.1. Boundary Conditions

So far, we have evaluated the characteristic function  $\varphi_Z = \varphi_Z(t)$  at specific values for  $t$ . Additional necessary conditions can be deduced by considering the asymptotic behavior of the characteristic function  $\varphi_Z(t)$  for  $t \rightarrow \infty$ . Intuitively, large values for  $t$  correspond to the case where the cosine term is changing sign very frequently, so that the integral approaches zero over intervals where  $g$  is smooth. And indeed, as discussed in Erdélyi [24], the asymptotics of  $\varphi_Z$  depend entirely on the behavior of the integrand in the neighborhood of certain distinguished points, called *critical points*. These critical points are either the endpoints of the interval of integration or the points at which the integrand (or some derivative thereof) exhibits a discontinuity. Given this intuition, it should not be too surprising that one may obtain the following auxiliary result for compactly supported DFD densities.



**Lemma 4.** Suppose that  $g : [-1, 1] \rightarrow \mathbb{R}_{\geq 0}$  is  $N$ -times continuously differentiable as well as DFD. Then,  $g^{(M)}(1) = g^{(M)}(-1) = 0$ , for any  $M \in \{0, \dots, N-1\}$ .<sup>13</sup>

**Proof.** See the appendix.  $\square$

## 4.2. Analyticity

A function is *analytic* if it admits derivatives of any finite order and can be extended into a Taylor series at each point of its domain of definition. Examples for analytic functions are polynomials and the exponential function. Sums, differences, and products of analytic functions are likewise analytic. On the other hand, the analyticity condition is violated, for example, when higher-order differentiability fails or when, even though derivatives of all orders exist, the Taylor series does not locally converge to the density function.

Letting  $N \rightarrow \infty$  in Lemma 4 and subsequently exploiting the fact that an analytic function is identified by its derivatives at any single point of its domain of definition, we arrive at the following useful observation.

**Proposition 4.** Suppose that  $g$  is the density of a compactly supported DFD distribution. Then,  $g$  is not equal on its support to some analytic function.

**Proof.** See the appendix.  $\square$

Proposition 4 may be used to extend our earlier observations regarding the beta distribution. Indeed, the beta density with integer parameter  $\alpha \in \mathbb{N} = \{1, 2, \dots\}$  is a polynomial on  $[-1, 1]$ , hence analytic. Therefore, the beta density is not DFD for any integer value  $\alpha > 0$ . In particular, this includes cases (viz., for  $\alpha = 3, 4, \dots$ ) where the density is neither convex nor concave. A similar example is the raised cosine, which likewise admits an analytic density and consequently is not DFD.

## 4.3. Kinks

As illustrated by the examples of the triangular and the Laplace distribution, DFD densities may exhibit a kink at the origin (cf. Figure 2). In fact, the triangular density, if considered as a function on the real line, has two additional kinks, namely, at the boundary of its support. The following result shows that such kinks are, under smoothness conditions on the component densities, a quite typical feature of a DFD density with bounded support.

**Proposition 5.** Let  $X$  and  $Y$  be i.i.d. random variables admitting a continuously differentiable density  $f$  on  $[0, 1]$ .

- (i) If  $\max\{f(0), f(1)\} > 0$ , then  $g$  has a kink at the origin.
- (ii) If  $\min\{f(0), f(1)\} > 0$ , then  $g$  has kinks also at  $\pm 1$ .

**Proof.** See the appendix.  $\square$

Thus, if the component density is positive at *at least one* boundary point of its support interval, then the density  $g$  of the DFD distribution necessarily exhibits a kink at the center of its support. If the component density is positive even at *both* boundary points of its support interval (as in the case of the uniform density), then  $g$  exhibits additional kinks at the boundary points of its own support interval.

## 5. Sufficient Conditions and the Construction of Components

This section explores conditions sufficient for a density to be DFD and derives a formula for the construction of the difference-form component under those conditions.

**Proposition 6.** Suppose that  $\varphi_Z \geq 0$  and that  $\sqrt{\varphi_Z}$  is positive definite. Then,  $Z$  is DFD, and a difference-form component of  $Z$  is given by the density function

$$f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \sqrt{\varphi_Z(t)} \cos(tx) dt. \quad (1)$$

**Proof.** See the appendix.  $\square$

Positive definiteness of the square root of  $\sqrt{\varphi_Z}$  intuitively imposes restrictions on the shape of  $\varphi_Z$ . We will discuss the positive definiteness condition in more detail further below. In principle, any of the sufficient conditions known for characteristic functions may be used to check the conditions of Proposition 6. For example, in a straightforward application of Pólya's condition, if  $\sqrt{\varphi_Z}$  is convex on  $\mathbb{R}_{\geq 0}$ , then  $Z$  is DFD. In applications, however, the often most convenient way to verify positive definiteness of a function is by checking that the inverse Fourier transform (1) is

globally nonnegative. For illustration of this approach, we reconstruct the uniform component from the triangular distribution.

**Example 3.** The characteristic function of the triangular density on  $[-1, 1]$  is given as  $\varphi_Z(t) = \frac{4 \sin^2(t/2)}{t^2} \geq 0$ . To find a difference-form component, we apply Equation (1). This yields

$$\begin{aligned} f(x) &= \frac{1}{\pi} \int_{-\infty}^{\infty} \sin\left(\frac{t}{2}\right) \cos(tx) \frac{dt}{t} \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \left\{ \sin\left(t\left(\frac{1}{2} + x\right)\right) + \sin\left(t\left(\frac{1}{2} - x\right)\right) \right\} \frac{dt}{t} \\ &= \frac{1}{\pi} \int_0^{\infty} \frac{\sin t}{t} dt \\ &= 1. \end{aligned}$$

Thus,  $\sqrt{\varphi_Z(t)}$  is positive definite, and by Proposition 6, the uniform distribution on the unit interval has been retrieved as a difference-form component of the triangular distribution.<sup>14</sup>

Analogous calculations are feasible, for instance, for the normal, Cauchy, and Laplace distributions. See Table 1 for details.

The proof of Proposition 6 is not deep but abstract. Technically, the assumptions of Proposition 6 ensure that  $g$  admits a convolution root that is a symmetric density. This automatically leads to a condition sufficient for difference-form decomposability, because any sum of two symmetric i.i.d. random variables is obviously DFD.<sup>15</sup>

Finally, recall that a characteristic function  $\varphi$  is called *infinitely divisible* if, for every positive integer  $n$ , there is a characteristic function  $\phi$  such that  $\varphi = \phi^n$ . A probability distribution is called infinitely divisible if its characteristic function is infinitely divisible (e.g., Lukacs [45, p. 19]).

**Corollary 1.** Any symmetric infinitely divisible distribution is DFD.

**Proof.** See the appendix.  $\square$

For instance, the generalized normal distribution where  $g(z)/g(0) = \exp\left(-\frac{|z|^p}{2}\right)$  with shape parameter  $p > 0$  (example kindly suggested by Felix Várdy) is DFD for  $0 < p \leq 1$  and  $p = 2$  because it is infinitely divisible (see Dytso et al. [21, theorem 5]). However, it is not DFD for  $p > 2$  because its characteristic function changes sign at least once in that case (see Dytso et al. [21, theorem 3]). For  $1 < p < 2$ , it is not known if this distribution is DFD.

## 6. Extensions


### 6.1. Distributions with Finite Support

One might wonder if a consideration of distributions with finite support might help to shed light on the class of DFD distributions. The insights from such exercise are limited, however. To understand why, consider the simplest case of an equidistant grid. Suppose we are given a vector of probabilities  $(p_0, p_1, \dots, p_N)$ , for  $N \geq 1$ , such that  $p_0 + 2\sum_{n=1}^N p_n = 1$ . The interpretation is that  $p_n$  corresponds to the probability that the symmetric random variable  $Z$  realizes as  $n \in \{0, \dots, N\}$ , and for any  $n > 0$ , likewise to the probability that  $Z$  realizes as  $-n$ . A difference-form component  $X$ , if it exists, may then be represented by a vector of probabilities  $(q_0, \dots, q_N)$ , where  $q_n$  denotes the probability that  $X$  realizes as  $n \in \{0, \dots, N\}$ . The system of equations to be solved is the following (cf. Hodges and Lehmann [34]):

$$\begin{aligned} p_0 &= q_0^2 + \dots + q_N^2 \\ p_1 &= q_0 q_1 + \dots + q_{N-1} q_N \\ &\vdots \\ p_{N-1} &= q_0 q_{N-1} + q_1 q_N \\ p_N &= q_0 q_N. \end{aligned} \tag{2}$$

The set of vectors  $(p_0, p_1, \dots, p_N)$  for which a solution  $(q_0, \dots, q_N)$  exists may, in principle, be characterized in explicit terms.

**Figure 5.** Illustration of the set of DFD distributions.



**Proposition 7.** For any fixed  $N \geq 2$ , the set of DFD discrete distributions forms a semialgebraic set, that is, it may be described by a finite number of algebraic identities and inequalities in the variables  $(p_0, p_1, \dots, p_N)$ .

**Proof.** See the appendix.  $\square$

We illustrate this general result with the help of a tractable example, the details of which may be found in the appendix.

**Example 4.** For  $N = 2$ , a distribution given by  $(p_0, p_1, p_2)$  is DFD if and only if  $p_1 \leq \frac{1}{4}$  and  $p_2 \leq \frac{(1+\sqrt{1-4p_1})^2}{16}$ . In this case, the set of difference-form components may be described in explicit terms.

Figure 5 illustrates the set of DFD distributions for  $N = 2$  as the area below the thick curve. The straight lines correspond to three nonnegativity constraints of the discrete cosine transform (defined in analogy to the continuous case), which are  $p_1 \leq \frac{1}{4}$ ,  $p_1 + p_2 \leq \frac{1}{3}$ , and  $p_1 + 2p_2 \leq \frac{1}{2}$ . As can be seen, the set of DFD distributions is a *strict* subset of the distributions with nonnegative cosine transform. For example,  $(p_0, p_1, p_2) = (\frac{1}{3}, \frac{1}{6}, \frac{1}{6})$  satisfies the nonnegativity constraints but is not DFD. This observation is in line with the corresponding fact for the continuous case, that is, that a nonnegative cosine transform is a necessary, but not a sufficient, condition for difference-form decomposability.

For  $N \geq 3$ , however, running the Tarski-Seidenberg algorithm that leads to a characterization of the set of DFD distributions through a finite number of algebraic identities and inequalities, although theoretically feasible, becomes substantially more involved. Moreover, determining a difference-form component in explicit form ceases to be tractable for  $N \geq 4$ .<sup>16</sup>

## 6.2. Functional Inequalities

Additional necessary conditions on the shape of a DFD density may be derived if the corresponding characteristic function is *integrable*. The following lemma provides a simple condition sufficient for this to be the case.


**Lemma 5.** Suppose that the difference-form component  $X$  of some random variable  $Z$  is distributed according to some continuously differentiable density function  $f_X$  with compact support. Then,  $\varphi_Z$  is integrable.

**Proof.** See the appendix.  $\square$

We are now all set to state our result regarding functional inequalities.

**Proposition 8.** Suppose that a DFD random variable  $Z$  is distributed according to some density function  $g$ . Assume also that  $\varphi_Z$  is integrable. Then,  $g$  is positive definite. In particular, the following inequalities hold:

- (i)  $g(z) \leq g(0)$  for any  $z \in \mathbb{R}$ ;
- (ii) if  $g(z) = 0$  outside of  $[-1, 1]$ , then  $g(z) \leq g(0) \cos\left(\frac{\pi}{1+|1/z|}\right)$  for any  $z \in (0, \frac{1}{2}]$ .<sup>17</sup>

**Figure 6.** Illustration of the inequality in Proposition 8(ii).

**Proof.** See the appendix.  $\square$

Thus, a DFD density with integrable characteristic function is positive definite, which is a fairly strong property. For example, it follows that any continuous DFD density assumes its maximum at the origin (even if the component is not unimodal). This observation is intuitively in line with the interpretation of  $g$  as an autocorrelation function (cf. Section 2) and admits a simple direct proof.<sup>18</sup> The less obvious inequality in part (ii) says that  $g(z)/g(0)$  remains weakly below the staircase function displayed in Figure 6. This inequality was shown to be an implication of positive definiteness by Boas and Kac [9]. It should be noted that Proposition 8 is in a sense equivalent to Lemma 2 and, hence, equally strong in its implications. Thus, the two results may be seen as complementary methods for testing for difference-form decomposability.

To prove that  $g$  is positive definite under the conditions of Proposition 8, one combines two powerful theorems in the literature on characteristic functions, Bochner's theorem and the Fourier inversion theorem. Bochner's theorem says that a function  $\phi$  is a characteristic function if and only if  $\phi$  is continuous, is positive definite, and satisfies  $\phi(0) = 1$ .<sup>19</sup> The proof then proceeds as follows. By the Fourier inversion theorem, we may reconstruct a density from its characteristic function. In the case of a symmetric density, however, the cosine transform of the characteristic function coincides (up to a constant factor) with the transform that generates the characteristic function from the density. We may therefore interpret  $\varphi_Z$ , provided it is integrable, as a density  $\hat{g}$  (after suitable normalization) of some "dual" random variable  $\hat{Z}$  and apply Bochner's theorem to  $\hat{Z}$  to derive necessary properties of its characteristic function  $\hat{\phi}$  that happens to coincide with  $g$  (again, up to a constant factor).

Proposition 8 may be used to obtain another partial result for the beta distribution in the leftover case where  $\alpha > 2$  (and  $\alpha$  not an integer). Indeed,

$$\left. \frac{\partial^2}{\partial z^2} \cos\left(\frac{\pi}{1+1/z}\right) \right|_{z=0} = -\pi^2,$$

$$\left. \frac{\partial^2}{\partial z^2} (1-z^2)^{\alpha-1} \right|_{z=0} = -2(\alpha-1).$$

Thus, for  $\alpha < \alpha^*$ , where  $\alpha^* = \frac{\pi^2}{2} + 1 \approx 5.93$ , making use of Lemma 5, the density of the beta distribution cannot be represented as the difference-form convolution of two i.i.d. continuously differentiable densities.<sup>20</sup>

### 6.3. Ratio-Form Decomposability

In analogy to the analysis of DFD random variables, one may ask which random variables may be represented as the ratio  $Z = X/Y$  of two i.i.d. random variables  $X$  and  $Y$ , following Huntington [35] and Curtiss [17]. Given our focus on applications in economic theory, we restrict attention to the case where the components assume positive values only. Then, however, it is *always* feasible to transform the argument of the distribution function using the logarithm. For example, any lognormal distribution may be expressed as the ratio of two i.i.d. lognormal distributions, etc. This general observation also has implications for contest success functions of the ratio-form, such as Tullock's [70]. Specifically, if idiosyncratic *multiplicative* noise is distributed according to the inverse exponential distribution (Fu and Lu [27], Jia [37]), then the density is given as  $f(x) = \alpha m x^{-(m+1)} \exp(-\alpha x^{-m})$  on  $[0, \infty)$ , where  $\alpha > 0$  and  $m > 0$  are

parameters. Transforming the corresponding distribution function  $F(x) = \exp(-\alpha x^{-m})$  via the exponential transform  $x = \exp(x)$  leads to the distribution function of the *additive* noise term, which is  $F(x) = \exp(-\alpha \exp(-mx))$ , that is, a Gumbel distribution (cf. Table 1). Therefore, the stochastic foundation of the Tullock contest is a direct consequence of the fact, familiar from McFadden [46], that the logistic distribution is the symmetrization of the Gumbel distribution.

More generally, the isomorphism between DFD random variables and *ratio-form decomposable* (RFD) random variables means that all the formulas and results above have a corresponding version in the theory of RFD densities. For instance, corresponding to Definition 2, the transformed symmetry condition on a positive random variable  $Z$  reads  $Z \stackrel{d}{=} 1/Z$ , where the distribution function  $G$  satisfies  $G(z) = 1 - G(1/z)$ . Consequently, this is a simple necessary condition for a positive random variable  $Z$  to be RFD is the existence of a density function  $g$  that satisfies  $g(z) = g(1/z)/z^2$ . For example, the ratio of two independent uniformly distributed random variables admits the density  $g(z) = 1/(2 \max\{1, z^2\})$ , which is ratio-form symmetric. However, the integral transform changes, with the Fourier transform (or characteristic function) being replaced by the Mellin transform (Epstein [23]).<sup>21</sup> Further, the crucial Lemma 2 above translates into a necessary condition for ratio-form decomposability, given by

$$\int_0^\infty g(z)\sqrt{z} \cos(t \ln z) dz \geq 0 \quad (t \in \mathbb{R}).$$

Thus, even though the conditions for RFD distributions may look different from those for DFD distributions, the transformation of each and every individual definition and result is ultimately mechanical. In fact, some necessary conditions are even invariant with respect to a log transformation of the argument, such as quasiconvexity, kinkedness, and analyticity of the density. However, other conditions such as concavity of the density are not invariant and may therefore lead to less intuitive characterizations. Similarly, we are not aware of a simple method that could be used to decompose an RFD probability distribution given by data on an equidistant grid.

#### 6.4. Correlated Noise Terms

Throughout the analysis, we have assumed that idiosyncratic noise terms are independently distributed. Indeed, the i.i.d. setting remains a common and natural benchmark that allows economists to study important first-order effects rather than having to worry about more intricate implications of correlation. However, it may be hard to find an application in which one can *not* argue that some degree of correlation may exist. For instance, because of macroeconomic fluctuations, revenue figures are likely to be positively correlated across sales representatives, as are key performance indicators across asset managers. On the other hand, if noise reflects shifts in preferences, as it may be the case for vertical differentiation, location choice, probabilistic voting, random utility, or paired comparisons, correlation across individual noise terms may be negative. Thus, the independence assumption is probably hardly ever necessarily satisfied, and some correlation seems to be the natural assumption in many applications.

However, departing from the independence assumption, the correspondence between the convolution operator on the one hand and the multiplication of characteristic functions on the other would break down. Given that this correspondence plays a crucial role in the derivation of most of our results, we actually have very little to say about the case with correlation. In fact, it is not clear how a theory with the possibility of correlation between noise terms could be developed in the first place. Relatedly, with correlation, decomposability would become a gradual concept, where constraints on the degree of correlation (measured in some way or other) would determine if a decomposition is feasible or not. Thus, dropping the independence assumption would affect the very nature of the analysis. For such reasons, an analysis of DFD densities with correlation must be left to future research.

On an intuitive level, however, independence does not seem crucial for our main results, in the sense that, even with limited correlation, difference-form decomposability should still lead to a lot of structure on the density function. For instance, we conjecture that an approximately uniform distribution cannot be represented as a difference of two identically distributed noise terms even if some limited degree of correlation is allowed for.

### 7. Concluding Remarks

In numerous economic models, uncertainty enters through a noise term that corresponds to the difference of two i.i.d. random variables. Our results allow deciding in many cases which distributions admit an i.i.d. difference-form decomposition and which do not. Thus, the analysis sheds light on the interesting but elusive class of DFD distributions.

In addition, our analysis shows that imposing distributional assumptions on the difference term is far from innocuous. Even intuitively plausible assumptions on the density of the difference term, such as approximate uniformity, quasiconvexity, strict concavity, or compact support combined with analyticity are *always* inconsistent,



and therefore entail the risk of ending up with incorrect economic conclusions. We pointed out, in particular, that the lack of decomposability of the uniform distribution, which is relevant in a variety of modeling frameworks, is actually a rather robust phenomenon.

To avoid the numerous pitfalls identified by the present studies, the applied economist has essentially three choices. First, if a specific distribution with compact support is desired, the triangular distribution, with its uniform components, seems to be the most natural assumption. If instead the uniform difference must be chosen (e.g., to ensure tractability), then this would require a rationale (like perfect negative correlation; see Meyer [47] or Bagnoli et al. [6]) as well as a discussion of how a change in the assumptions about the distribution of noise would likely affect the results. Second, if a specific distribution with full support is desired, then any infinitely divisible distribution, with components taken from the same family, will do the job. Common examples are the normal and the Cauchy distribution families. Finally, if the researcher aims at keeping distributional assumptions at a minimum, then properties automatically fulfilled, like symmetry, positive definiteness, and functional inequalities, should be used. Any additional assumptions, such as unimodality (Hodges and Lehmann [34]) or logconcavity (Ibragimov [36]), should be imposed on the components, and the fact that such properties are inherited should be used to obtain conclusions on the difference term.

There are several dimensions in which the present study could be extended. Of interest, for instance, might be the consideration of multivariate distributions, that is, random variables with values in Banach spaces. More interesting, albeit also more challenging, might be the question of what happens if the number  $N$  of i.i.d. individual random terms is larger than two. In that case, one would have to study the joint distribution of  $\binom{N}{2}$  correlated difference terms, each of which would be DFD.

Finally, as noted by one of the reviewers, our properties and conditions might also prove useful in testing whether a given model with i.i.d. noise terms could fit observed data, especially for a setting with symmetric players and observable inputs. In that case, the test would concern the properties of the difference of random indicators of different options.<sup>22</sup>

## Acknowledgments

The authors thank the handling editor and two anonymous reviewers for the efficient review process. For useful comments and discussions, the authors thank Kai Konrad and Damian Kozbur, as well as participants of the SAET Conference in Paris in July 2023.

## Appendix. Proofs

This appendix contains technical material omitted from the body of the paper.

**Details on Example 1.** Suppose that  $Z \sim \mathcal{U}[-1, 1]$ . The characteristic function of  $Z$  is

$$\varphi_Z(t) = \frac{1}{2} \int_{-1}^1 \cos(tz) dz = \frac{\sin t}{t}.$$

It is then clear that  $\varphi_Z(t) < 0$  for selected values of  $t$ , in conflict with Lemma 2.  $\square$

**Proof of Proposition 1.** Evaluating the characteristic function at  $t = \frac{3\pi}{2}$  yields

$$\begin{aligned} \varphi_Z\left(\frac{3\pi}{2}\right) &= \int_{-1/3}^{1/3} \underbrace{\cos\left(\frac{3\pi}{2}x\right)}_{\geq 0} g(x) dx + 2 \int_{1/3}^1 \underbrace{\cos\left(\frac{3\pi}{2}x\right)}_{\leq 0} g(x) dx \\ &< \int_{-1/3}^{1/3} \cos\left(\frac{3\pi}{2}x\right) \left(\frac{1}{2} + \frac{1}{6}\right) dx + 2 \int_{1/3}^1 \cos\left(\frac{3\pi}{2}x\right) \left(\frac{1}{2} - \frac{1}{6}\right) dx \\ &= 0. \end{aligned}$$

This proves the claim.  $\square$

**Details on Example 2.** On the unit interval, the density of the beta distribution with shape parameters  $\alpha > 0, \beta > 0$  is commonly defined as

$$g_{[0,1]}(x) = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha-1} (1-x)^{\beta-1} \quad (x \in [0, 1]).$$

A stretched variant with support  $[-1, 1]$  is given as

$$g(z) \equiv \frac{1}{2} g_{[0,1]} \left( \frac{z+1}{2} \right) = \frac{\Gamma(\alpha+\beta)}{2^{\alpha+\beta-1} \Gamma(\alpha) \Gamma(\beta)} (1+z)^{\alpha-1} (1-z)^{\beta-1}.$$

Assuming  $\alpha = \beta$ , and exploiting the duplication rule for the gamma function, that is,  $\Gamma(2\alpha) = 2^{2\alpha-1} \pi^{-1/2} \Gamma(\alpha) \Gamma(\alpha + \frac{1}{2})$ , leads to the parametric form considered in the body of the paper. The shape of the beta density is determined by its second derivative

$$\frac{\partial^2 (1-x^2)^{\alpha-1}}{\partial x^2} = 2(1-\alpha)(1-x^2)^{\alpha-3} (2x^2(2-\alpha) + (1-x^2)).$$

Hence, the beta density  $g$  is convex for  $\alpha \in (0, 1]$ , strictly concave for  $\alpha \in (1, 2]$ , and neither convex nor concave for  $\alpha > 2$ .

**Proof of Proposition 2.** By contradiction. Suppose that  $Z$  is DFD. Then, by Lemma 2,  $\varphi_Z(t) \geq 0$  for any  $t > 0$ . However, evaluating  $\varphi_Z(t)$  at  $t = 2\pi$ , we see that

$$0 \leq \varphi_Z(2\pi) = \int_{-1}^1 \cos(2\pi z) g(z) dz = \frac{1}{2\pi} \int_0^{2\pi} \cos(\hat{z}) g\left(\frac{\hat{z}}{2\pi}\right) d\hat{z},$$

with  $\hat{z} = 2\pi z$ . Because  $\cos(\hat{z} + \pi) = -\cos(\hat{z})$ , it follows that

$$\begin{aligned} \varphi_Z(2\pi) &= \frac{1}{2\pi} \int_0^\pi \cos(\hat{z}) g\left(\frac{\hat{z}}{2\pi}\right) d\hat{z} + \cos(\hat{z} + \pi) g\left(\frac{\hat{z} + \pi}{2\pi}\right) d\hat{z} \\ &= \frac{1}{2\pi} \int_0^\pi \cos(\hat{z}) \left\{ g\left(\frac{\hat{z}}{2\pi}\right) - g\left(\frac{\hat{z} + \pi}{2\pi}\right) \right\} d\hat{z} \\ &< 0, \end{aligned}$$

where the inequality is strict because  $g$  is not uniform by Example 1. The contradiction proves the claim.  $\square$

**Proof of Proposition 3.** Suppose that  $g(z)$  is DFD and strictly concave on its support. Then,  $g(z)$  is symmetric by Lemma 1. Clearly, therefore,  $Z$  has compact support, say  $[-1, 1]$ . Moreover, by standard results on concave functions (e.g., Royden and Fitzpatrick [62, p. 117]), the derivative  $g'$  is well defined except possibly at kinks that form a set of measure zero. Moreover,  $g'$  is strictly declining. Hence, evaluating the integral transform at  $t = 2\pi$ , integration by parts delivers

$$\begin{aligned} \varphi(2\pi) &= \int_{-1}^1 \cos(2\pi z) g(z) dz \\ &= \underbrace{\frac{1}{2\pi} \sin(2\pi z) g(z) \Big|_{-1}^1}_{=0} - \frac{1}{\pi} \int_0^1 \sin(2\pi z) g'(z) dz \\ &< \frac{1}{\pi} \int_0^{1/2} \underbrace{\left( \sin(2\pi z) + \sin\left(2\pi\left(z + \frac{1}{2}\right)\right) \right)}_{=0} g'(z) dz, \end{aligned}$$

but this is in conflict with Lemma 2.  $\square$

The following auxiliary result will be used in the proofs of Lemmas 4 and 5. As usual,  $o(t^{-N})$  denotes a function that goes to zero more quickly than  $t^{-N}$  (i.e.,  $\lim_{t \rightarrow \infty} t^N o(t^{-N}) = 0$ ).

**Lemma A.1** (A. Erdélyi). *Suppose that  $g(z)$  is  $N$ -times continuously differentiable on the interval  $[\alpha, \beta]$ , where  $-\infty < \alpha < \beta < \infty$ . Then,*

$$\int_\alpha^\beta g(z) \exp(itz) dz = \Phi_N(t, \beta) - \Phi_N(t, \alpha) + o(t^{-N}),$$

where  $i = \sqrt{-1}$ , and

$$\Phi_N(t, z) = \sum_{M=0}^{N-1} i^{M-1} g^{(M)}(z) \frac{\exp(itz)}{t^{M+1}}.$$

**Proof.** See Erdélyi [24, theorem 1].  $\square$

**Proof of Lemma 4.** By induction. (*Induction basis*) Suppose that  $g$  is continuously differentiable on  $[-1, 1]$  as well as DFD. Because  $g$  is continuously differentiable,  $\Phi_1(t, 1) = g(1) \frac{\sin(t)}{t} = -\Phi_1(t, -1)$ . Moreover, because  $g$  is DFD, Lemmas 2 and A.1 jointly imply that

$$0 \leq \varphi_Z(t) = \int_1^1 g(z) \cos(zt) dz = 2g(1) \frac{\sin t}{t} + o\left(\frac{1}{t}\right).$$

For  $t$  large, the  $\frac{\sin t}{t}$  term dominates, so that necessarily  $g(1) = g(-1) = 0$ . (*Induction step*) Let  $N \geq 1$ , and assume that the claim has been shown for  $N$ . Suppose that  $g$  is  $(N+1)$ -times continuously differentiable. Then,  $g$  is  $N$ -times continuously differentiable so that, from the induction hypothesis,  $g^{(M)}(1) = g^{(M)}(-1) = 0$  for any  $M \in \{0, \dots, N-1\}$ . Now, from the definition of  $\Phi_{N+1}(t, 1)$  and  $\exp(it) = \cos(t) + i \sin(t)$ ,

$$\Phi_{N+1}(t, 1) = \sin(z) \left( \sum_{k=0}^{\lfloor \frac{N}{2} \rfloor} \frac{(-1)^k}{t^{2k+1}} g^{(2k)}(1) \right) + \cos(z) \left( \sum_{k=1}^{\lfloor \frac{N+1}{2} \rfloor} \frac{(-1)^{k+1}}{t^{2k}} g^{(2k-1)}(1) \right).$$

If  $N = 2K$  is even, then

$$\Phi_{N+1}(t, 1) = \frac{(-1)^K \sin(z)}{t^{N+1}} g^{(N)}(1) = -\Phi_{N+1}(t, -1),$$

so that  $g^{(N)}(1) = g^{(N)}(-1) = 0$ . If  $N = 2K - 1$  is odd, then

$$\Phi_{N+1}(t, 1) = \frac{(-1)^{K+1} \cos(z)}{t^{N+1}} g^{(N)}(1) = -\Phi_{N+1}(t, -1),$$

and we find  $g^{(N)}(1) = g^{(N)}(-1) = 0$ , as before. This proves the claim.  $\square$

**Proof of Proposition 4.** Suppose that  $g$  is DFD. Because  $g$  is analytic at  $-1$  and at  $1$ , it is infinitely differentiable there, so that all derivatives at  $-1$  and at  $1$  are zero by Lemma 4. By the identity theorem for analytic functions, this implies that  $g$  vanishes on  $[-1, 1]$ , which is impossible. The contradiction shows that  $g$  cannot be DFD.  $\square$

**Proof of Proposition 5.** For  $z \in [0, 1]$ , we have that

$$g(z) = \int_0^{1-z} f(z+x)f(x)dx.$$

Hence, using Leibniz's rule,

$$\begin{aligned} \lim_{z \searrow 0} \frac{g(z) - g(0)}{z} &= \lim_{z \searrow 0} \frac{1}{z} \left\{ \int_0^{1-z} f(z+x)f(x)dx - \int_0^1 f(z+x)f(x)dx \right\} \\ &= -\lim_{z \searrow 0} \frac{1}{z} \int_{1-z}^1 f(z+x)f(x)dx \\ &= -f(1)^2 + \int_0^1 f'(x)f(x)dx \\ &= -f(1)^2 + \frac{f(1)^2 - f(0)^2}{2} \\ &= -\frac{f(0)^2 + f(1)^2}{2} \\ &< 0. \end{aligned}$$

On the other hand, by symmetry,  $g(z) = g(-z)$ , so that

$$\lim_{z \nearrow 0} \frac{g(z) - g(0)}{z} > 0.$$

Similarly, one finds

$$\lim_{z \nearrow 1} \frac{g(z) - g(1)}{z} = \lim_{z \nearrow 1} \frac{1}{z} \left\{ \int_0^{1-z} f(z+x)f(x)dx \right\} = -f(1)f(0),$$

and a corresponding expression at  $z = -1$ . This proves the proposition.  $\square$

**Proof of Proposition 6.** Immediate from Bochner’s theorem and the Fourier inversion theorem.  $\square$

**Proof of Corollary 1.** Let the distribution of  $Z$  be infinitely divisible and symmetric. Then,  $\varphi_Z \geq 0$  is continuous (by Bochner’s theorem), and does not possess any real zeros (Lukacs [45]). Let  $\phi$  be a characteristic function such that  $\varphi = \phi^2$ . Then,  $\phi > 0$  or  $\phi < 0$ , but the second alternative is not feasible because  $\phi(0) = 1$ . Hence,  $\phi > 0$ , and  $\varphi = |\phi|^2$ . The claim follows.  $\square$

**Proof of Proposition 7.** Immediate from the Tarski-Seidenberg theorem (e.g., Neyman and Sorin [52, p. 65]).  $\square$

**Details on Example 4.** Suppose that  $N=2$  and fix a vector of probabilities  $(p_0, p_1, p_2)$  such that  $p_0 + 2p_1 + 2p_2 = 1$ . Taking account of redundancies, a vector of probabilities  $(q_0, q_1, q_2)$  is sought such that

$$p_1 = q_0 q_1 + q_1 q_2 \quad (\text{A.1})$$

$$p_2 = q_0 q_2 \quad (\text{A.2})$$

$$1 = q_0 + q_1 + q_2. \quad (\text{A.3})$$

Combining (A.1) and (A.3), we see that  $p_1 = q_1(1 - q_1)$ , hence necessarily  $p_1 \leq \frac{1}{4}$  and  $q_1 \in \{q_1^+, q_1^-\}$ , where  $q_1^+ = \frac{1}{2} + \sqrt{\frac{1}{4} - p_1}$  and  $q_1^- = \frac{1}{2} - \sqrt{\frac{1}{4} - p_1}$ . From (A.2) and (A.3), one notes that  $q_0 + q_2 = 1 - q_1$  and  $q_0 q_2 = p_2$ . Therefore, a difference-form component with  $q_1 \in \{q_1^+, q_1^-\}$  exists if and only if  $4p_2 \leq (1 - q_1)^2$ , and the corresponding solution, or pair of solutions, is given by

$$\{q_0, q_2\} = \frac{1 - q_1}{2} \pm \sqrt{\frac{(1 - q_1)^2}{4} - p_2}.$$

Next, we note that  $q_1^- \leq q_1^+ \leq 1$ , so that  $(1 - q_1^-)^2 \geq (1 - q_1^+)^2$ . Thus, a necessary and sufficient condition for a solution to exist is  $p_1 \leq \frac{1}{4}$  and  $p_2 \leq \frac{1}{16}(1 + \sqrt{1 - 4p_1})^2$ , as claimed. Moreover, there are at most four solutions.<sup>23</sup>

**Proof of Lemma 5.** Because  $f$  is continuous differentiable, Lemma A.1 implies

$$\varphi_X(t) = \int_0^c f(z) \exp(izt) dz = \frac{1}{t} (\sin(ct) + i(f(0) - \cos(ct)f(c))) + o(t^{-1}).$$

By Lemma 2,

$$\varphi_Z(t) = |\varphi_X(t)|^2 = \frac{(\sin(ct)f(c))^2 + (f(0) - \cos(ct)f(c))^2}{t^2} + o(t^{-2}).$$

Thus,  $\varphi_Z$  is integrable, as claimed.  $\square$

**Proof of Proposition 8.** Suppose that  $Z$  is DFD. Then, by Lemma 2,  $\varphi_Z(t) \geq 0$  for all  $t \in \mathbb{R}$ . Moreover,  $\varphi_Z$  is continuous by Bochner’s theorem, with  $\varphi_Z(0) = \int_{-\infty}^{\infty} g(z) dz = 1$ . Hence, using that  $\varphi_Z$  is integrable, we see that  $\int_{-\infty}^{\infty} \varphi_Z(s) ds > 0$ . Moreover, by the Fourier inversion theorem,  $g(0) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \varphi_Z(s) ds$ . Let

$$\hat{g}(z) = \frac{\varphi_Z(z)}{\int_{-\infty}^{\infty} \varphi_Z(s) ds} = \frac{\varphi_Z(z)}{2\pi g(0)}.$$

Then,  $\hat{g}$  is a density of some random variable  $\hat{Z}$ . The characteristic function of  $\hat{Z}$  is, therefore, given by

$$\varphi_{\hat{Z}}(t) = \frac{1}{g(0)} \left( \frac{1}{2\pi} \int_{-\infty}^{\infty} \varphi_Z(z) \cos(tz) dt \right) = \frac{g(t)}{g(0)},$$

where we applied again the Fourier inversion theorem. But, by Bochner’s theorem, the characteristic function of  $\hat{Z}$  is positive definite. Hence,  $g$  is positive definite as well. Inequality (i) is an immediate consequence of the definition of positive definiteness. Inequality (ii) was shown to follow from positive definiteness by Boas and Kac [9].  $\square$

## Endnotes

<sup>1</sup> The related literature will be reviewed later in this section.

<sup>2</sup> Lazear and Rosen [38, p. 860] wrote “Since  $g$  is symmetric and nonuniform [ ... ],” where  $g$  is the distribution of the difference between the two (worker-idiosyncratic) i.i.d. random components. See also Drugov and Ryvkin [19, footnote 3].

<sup>3</sup> Several recent contributions have stressed the salience of the shape of noise for economic predictions (e.g., Drugov and Ryvkin [18], Drugov and Ryvkin [19], Morgan et al. [50], Morgan et al. [51]).

<sup>4</sup> Our sufficient conditions loosely relate to the *Wiener-Khinchine-Kolmogorov criterion* and *convolution roots* studied by Boas and Kac [9]. See also Ehm et al. [22] and Akopyan and Efimov [2]. However, although that literature admits such roots to be complex-valued, we are seeking convolution roots that are probability densities, that is, that are real-valued and nonnegative.

<sup>5</sup> Note that we do not take the absolute value of the difference. Puri and Rubin [57] and Stadje [65] studied distributions with the property that the absolute difference  $Z = |X - Y|$  is identically distributed as its two i.i.d. components  $X$  and  $Y$ . Interestingly, if the distribution admits a density, then this property characterizes the exponential distribution.

<sup>6</sup> It is immediate to see that if  $X$  is a difference-form component of  $Z$ , then so is  $-X$ . In fact, this is even the case for  $c + X$  and  $c - X$ , for any constant  $c \in \mathbb{R}$ . Thus, any difference-form decomposition, provided it exists, can be unique at most up to reflection at the origin and arbitrary translations.

<sup>7</sup> The moment-generating function  $E[e^{tZ}]$  has the same advantage of being real-valued, but its use would obscure the idea of the frequency analysis conducted below. In addition, the moment-generating function does not exist for all distributions of interest (e.g., for the Cauchy distribution).

<sup>8</sup> The fact that the difference of two i.i.d. uniform random terms follows a triangular density has found numerous applications. See, e.g., Bull et al. [11], Schotter and Weigelt [64], Prendergast [56], Altmann et al. [4], Moldovanu et al. [48], and Ewerhart [25].

<sup>9</sup> Lewis [39] characterized the complete set of decompositions of the uniform distribution into arbitrarily many independent, but not necessarily identically distributed, components. See also Tortrat [68], cited by Ruzsa [63], and Topolyan [67].

<sup>10</sup> Further intuition may be gained from considering a discrete setting (see also Section 6). Suppose that the components  $X$  and  $Y$  independently realize as  $x = 0$  with probability  $q_0$  and as  $x = 1$  with probability  $q_1$ . Then,  $Z = X - Y$  realizes as  $z = 0$  with probability  $p_0 = q_0^2 + q_1^2$ , and as  $z = 1$  with probability  $p_1 = q_0q_1$ . Hence,  $p_0 - 2p_1 \geq 0$  if  $Z$  is DFD. Thus,  $p_0 \geq \frac{1}{2}$  and  $p_1 \leq \frac{1}{4}$ , that is, the uniform distribution on  $\{-1, 0, 1\}$  is not DFD.

<sup>11</sup> See Figure 2 for illustration. Details on this example can be found in the appendix.

<sup>12</sup> Pólya's sufficient criterion says that a function  $\phi$  is a characteristic function if  $\phi$  is continuous, is convex on the positive real line, and satisfies  $\phi(0) = 1$ . For an account of Pólya's sufficient condition and other sufficient conditions, we refer the reader to Lukacs' [45] survey.

<sup>13</sup> As usual,  $g^{(M)}(z)$  denotes the  $M$ -th derivative of  $g$  at  $z$ , provided it exists. In particular,  $g^{(0)}(z) = g(z)$ .

<sup>14</sup> In fact, this decomposition is unique (cf. O'Neill and Walther [53]).

<sup>15</sup> Notably, the converse statement is not generally true. That is, there are DFD densities that do not admit a symmetric difference-form component. For example, the component  $X$  given by the density  $f(x) = 2x$  on  $[0, 1]$  cannot be replaced by any symmetric component (Carnal and Dozzi [12, p. 168]). Relatedly, Gushchin and Küchler [32] have shown that any weakly declining density on  $\mathbb{R}_{\geq 0}$  is a.e. uniquely determined by its symmetrization.

<sup>16</sup> However, in analogy to Proposition 6, a *symmetric* difference-form component (i.e., a component that satisfies  $q_n = q_{N-n}$  for all  $n \in \{0, \dots, \lceil \frac{N}{2} \rceil - 1\}$ ), may always be computed (provided it exists) from system (2).

<sup>17</sup> Here,  $\lfloor 1/z \rfloor$  denotes the largest integer weakly smaller than the ratio  $1/z$ .

<sup>18</sup> Specifically, it suffices to note that  $g(0) - g(z) = \frac{1}{2} \int_{-\infty}^{\infty} (f(x+z) - f(x))^2 dx \geq 0$ .

<sup>19</sup> A real-valued function  $\phi$  is *positive definite* if, for every  $n \in \{1, 2, \dots\}$  and  $x_1, \dots, x_n \in \mathbb{R}$ , the matrix  $[\phi(x_i - x_j)]_{i,j=1}^n$  is positive semidefinite.

<sup>20</sup> The 123 Theorem (Alon and Yuster [3]), suitably reformulated, says that if  $G$  is the distribution function of a DFD random variable, then  $G(b) - G(a) \leq 2(\lfloor b/a \rfloor - 1)(G(a) - G(0))$ , for any  $b > a > 0$ , where  $\lfloor b/a \rfloor$  denotes the lowest integer weakly larger than the ratio  $b/a$ . That result, however, cannot be used to easily derive Proposition 8 or any other result of the present paper.

<sup>21</sup> As a result, the density function  $g$  of an RFD random variable may be expressed as

$$g(z) = \frac{1}{z^2} \int_0^{\infty} xf(x)f(x/z)dx,$$

where  $f$  denotes the density of the ratio-form component.

<sup>22</sup> A complication is that typically the data are not from a fixed pair of players, for example, in a contest setting, even when we assume for simplicity that all these players are symmetric. Such questions of interest, like others that remained unanswered in the present study, might, however, suggest worthwhile avenues for future research.

<sup>23</sup> For the special case  $N = 2$ , this confirms a conjecture of Carnal and Dozzi [12, p. 172], according to which the number of difference-form decompositions is at most  $2^N$ .

## References

- [1] Ahmed N, Natarajan T, Rao KR (1974) Discrete cosine transform. *IEEE Trans. Comput.* 100(1):90–93.
- [2] Akopyan R, Efimov A (2017) Boas–Kac roots of positive definite functions of several variables. *Anal. Math.* 43(3):359–369.
- [3] Alon N, Yuster R (1995) The 123 Theorem and its extensions. *J. Combin. Theory Ser. A* 72(2):322–331.
- [4] Altmann S, Falk A, Wibral M (2012) Promotions and incentives: The case of multistage elimination tournaments. *J. Labor Econom.* 30(1):149–174.
- [5] An MY (1998) Logconcavity vs. logconvexity: A complete characterization. *J. Econom. Theory* 80(2):350–369.
- [6] Bagnoli M, Viswanathan S, Holden C (2001) On the existence of linear equilibria in models of market making. *Math. Finance* 11(1):1–31.
- [7] Barlevy G, Neal D (2012) Pay for percentile. *Amer. Econom. Rev.* 102(5):1805–1831.
- [8] Becker GM, DeGroot MH, Marschak J (1963) Stochastic models of choice behavior. *Behav. Sci.* 8(1):41–55.
- [9] Boas R Jr, Kac M (1945) Inequalities for Fourier transforms of positive functions. *Duke Math. J.* 12(1):189–206.



- [10] Bradley RA, Terry ME (1952) Rank analysis of incomplete block designs: I. The method of paired comparisons. *Biometrika* 39(3/4): 324–345.
- [11] Bull C, Schotter A, Weigelt K (1987) Tournaments and piece rates: An experimental study. *J. Political Econom.* 95(1):1–33.
- [12] Carnal H, Dozzi M (1989) On a decomposition problem for multivariate probability measures. *J. Multivariate Anal.* 31(2):165–177.
- [13] Carson JR (1923) Method and means for signaling with high-frequency waves. U.S. Patent 1,449,382. <https://patents.google.com/patent/US1449382A/en>.
- [14] Che YK, Gale I (2000) Difference-form contests and the robustness of all-pay auctions. *Games Econom. Behav.* 30(1):22–43.
- [15] Chung KL (1953) Sur les lois de probabilité unimodales. *C. R. Math. Acad. Sci. Paris* 236(6):583–584.
- [16] Cramér H (1936) Über eine Eigenschaft der normalen Verteilungsfunktion. *Math. Z.* 41(1):405–414.
- [17] Curtiss JH (1941) On the distribution of the quotient of two chance variables. *Ann. Math. Statist.* 12(4):409–421.
- [18] Drugov M, Ryvkin D (2020) How noise affects effort in tournaments. *J. Econom. Theory* 188:105065.
- [19] Drugov M, Ryvkin D (2022) Hunting for the discouragement effect in contests. *Rev. Econom. Design.*, ePub ahead of print August 2, <https://doi.org/10.1007/s10058-022-00308-4>.
- [20] Dugué D (1957) *Arithmétique des lois de probabilités. Mémoires des Sciences Mathématiques* (Gauthier Villars, Paris).
- [21] Dytso A, Bustin R, Poor HV, Shamai S (2018) Analytical properties of generalized Gaussian distributions. *J. Statist. Distrib. Appl.* 5(1):1–40.
- [22] Ehm W, Gneiting T, Richards D (2004) Convolution roots of radial positive definite functions with compact support. *Trans. Amer. Math. Soc.* 356(11):4655–4685.
- [23] Epstein B (1948) Some applications of the Mellin transform in statistics. *Ann. Math. Statist.* 19(3):370–379.
- [24] Erdélyi A (1955) Asymptotic representations of Fourier integrals and the method of stationary phase. *J. Soc. Indust. Appl. Math.* 3(1):17–27.
- [25] Ewerhart C (2016) An envelope approach to tournament design. *J. Math. Econom.* 63:1–9.
- [26] Feller W (1971) *An Introduction to Probability Theory and Its Applications II*, 2nd ed. (Wiley, New York).
- [27] Fu Q, Lu J (2012) Micro foundations of multi-prize lottery contests: A perspective of noisy performance ranking. *Soc. Choice Welfare* 38(3):497–517.
- [28] Giraud BG, Peschanski R (2006) On positive functions with positive Fourier transforms. *Acta Phys. Polon. B* 37(2):331–346.
- [29] Glaeser EL, Ponzetto GA, Shapiro JM (2005) Strategic extremism: Why Republicans and Democrats divide on religious values. *Quart. J. Econom.* 120(4):1283–1330.
- [30] Goeree JK, Holt CA, Palfrey TR (2005) Regular quantal response equilibrium. *Experiment. Econom.* 8:347–367.
- [31] Gori F (2017) Doubly positive functions in coherent and partially coherent optics. *Optics Lett.* 42(23):4998–5001.
- [32] Gushchin AA, Kuchler U (2005) On recovery of a measure from its symmetrization. *Theory Probab. Appl.* 49(2):323–333.
- [33] Hirshleifer J (1989) Conflict and rent-seeking success functions: Ratio vs. difference models of relative success. *Public Choice* 63(2):101–112.
- [34] Hodges JL, Lehmann EL (1954) Matching in paired comparisons. *Ann. Math. Statist.* 25(4):787–791.
- [35] Huntington EV (1939) Frequency distribution of product and quotient. *Ann. Math. Statist.* 10(2):195–198.
- [36] Ibragimov IA (1956) On the composition of unimodal distributions. *Theory Probab. Appl.* 1(2):255–260.
- [37] Jia H (2008) A stochastic derivation of the ratio form of contest success functions. *Public Choice* 135(3/4):125–130.
- [38] Lazear EP, Rosen S (1981) Rank-order tournaments as optimum labor contracts. *J. Political Econom.* 89(5):841–864.
- [39] Lewis T (1967) The factorisation of the rectangular distribution. *J. Appl. Probab.* 4(3):529–542.
- [40] Lin YJ (1988) Oligopoly and vertical integration: Note. *Amer. Econom. Rev.* 78(1):251–254.
- [41] Lindbeck A, Weibull JW (1987) Balanced-budget redistribution as the outcome of political competition. *Public Choice* 52(3):273–297.
- [42] Linnik JV (1960) *Decomposition of Probability Distributions* (Oliver and Boyd, Edinburgh, UK).
- [43] Linnik JV, Ostrovskii IV (1977) *Decomposition of Random Variables and Vectors*. Translations of Mathematical Monographs, vol. 48 (American Mathematical Society, Providence, RI).
- [44] Lukacs E (1970) *Characteristic Functions*, 2nd ed. (Griffin, London).
- [45] Lukacs E (1972) A survey of the theory of characteristic functions. *Adv. Appl. Probab.* 4(1):1–37.
- [46] McFadden D (1973) Conditional logit analysis of qualitative choice behavior. Zarembka P, ed. *Frontiers in Econometrics* (Academic, New York), 105–142.
- [47] Meyer MA (1991) Learning from coarse information: Biased contests and career profiles. *Rev. Econom. Stud.* 58(1):15–41.
- [48] Moldovanu B, Sela A, Shi X (2012) Carrots and sticks: Prizes and punishments in contests. *Econom. Inquiry* 50(2):453–462.
- [49] Moretti E (2011) Local labor markets. Ashenfelter O, Card D, eds. *Handbook of Labor Economics*, vol. 4b (Elsevier, Amsterdam), 1237–1313.
- [50] Morgan J, Sisak D, Várdy F (2018) The ponds dilemma. *Econom. J. (London)* 128(611):1634–1682.
- [51] Morgan J, Tumlinson J, Várdy F (2022) The limits of meritocracy. *J. Econom. Theory* 201:105414.
- [52] Neyman A, Sorin S (2003) *Stochastic Games and Applications*. NATO Science Series C, vol. 570 (Springer, New York).
- [53] O’Neill EL, Walther A (1963) The question of phase in image formation. *Optica Acta (London)* 10(1):33–39.
- [54] Patterson AL (1935) A direct method for the determination of the components of interatomic distances in crystals. *Z. Kristallogr.* 90(1–6):517–542.
- [55] Pólya G (1949) Remarks on characteristic functions. Jerzy N, ed. *Proc. First Berkeley Conf. Math. Statist. Probab.* (University of California Press, Berkeley, CA), 115–123.
- [56] Prendergast C (2002) The tenuous trade-off between risk and incentives. *J. Political Econom.* 110(5):1071–1102.
- [57] Puri PS, Rubin H (1970) A characterization based on the absolute difference of two iid random variables. *Ann. Math. Statist.* 41(6): 2113–2122.
- [58] Puri ML, Sen PK (1968) On Chernoff-Savage tests for ordered alternatives in randomized blocks. *Ann. Math. Statist.* 39(3):967–972.
- [59] Roback J (1982) Wages, rents, and the quality of life. *J. Political Econom.* 90(6):1257–1278.
- [60] Rosen S (1979) Wage-based indexes of urban quality of life. Miezkowski PN, Straszheim MR, eds. *Current Issues in Urban Economics* (Johns Hopkins University Press, Baltimore), 74–104.
- [61] Rosenblatt J (1984) Phase retrieval. *Comm. Math. Phys.* 95(3):317–343.
- [62] Royden HL, Fitzpatrick P (1988) *Real Analysis*, 3rd ed. (Macmillan, New York).

- [63] Ruzsa IZ (1982) Arithmetic of probability distributions. *Séminaires Théor. Nombres Bordeaux* 20:1–12.
- [64] Schotter A, Weigelt K (1992) Asymmetric tournaments, equal opportunity laws, and affirmative action: Some experimental results. *Quart. J. Econom.* 107(2):511–539.
- [65] Stadje W (1994) A characterization of the exponential distribution involving absolute differences of iid random variables. *Proc. Amer. Math. Soc.* 121(1):237–243.
- [66] Thurstone LL (1927) A law of comparative judgment. *Psych. Rev.* 34(4):273–286.
- [67] Topolyan I (2014) Rent-seeking for a public good with additive contributions. *Soc. Choice Welfare* 42(2):465–476.
- [68] Tortrat A (1969) Sur un théorème de Lewis et la décomposition en facteurs premiers de la loi rectangulaire. *J. Appl. Probab.* 6(1):177–185.
- [69] Tuck EO (2006) On positivity of Fourier transforms. *Bull. Aust. Math. Soc.* 74(1):133–138.
- [70] Tullock G (1980) Efficient rent seeking. Buchanan JM, Tollison RD, Tullock G, eds. *Toward a Theory of the Rent-Seeking Society* (Texas A&M University Press, College Station, TX), 97–112.
- [71] Vogt H (1983) Unimodality of differences. *Metrika* 30(1):165–170.
- [72] Walther A (1963) The question of phase retrieval in optics. *Optica Acta (London)* 10(1):41–49.